ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Climate Change and Arboviral Diseases: Challenges for Surveillance and Primary Health Care

Maria Terezinha Gomes Pontes
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Edylene Maria dos Santos Pereira
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Bruno Ramos Brum
Doutorado em Ciencias Ambientais
Uninassau - Vilhena

Maria Aparecida de Almeida Araujo
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Thaita Thaisi Zago
Mestranda em Ciências Políticas
Universidad Europea del Atlántico, Espanha

Sérgio Raimundo Ernesto Machado
Doutorando em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Sheila Érika Ferro Ramalho Nobre
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Neide Moreira de Souza
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Abstract

Climate change has emerged as one of the greatest public health challenges of the twenty-first century, profoundly influencing the epidemiology of infectious diseases worldwide. Among the health conditions most directly affected by environmental transformations are arboviral diseases, including dengue, Zika, chikungunya, and yellow fever, whose transmission dynamics are highly sensitive to variations in temperature, rainfall, humidity, urbanization, and ecosystem disruption. These climatic and environmental changes have expanded the geographic distribution of mosquito vectors, prolonged transmission seasons, and increased the frequency and intensity of outbreaks in tropical and subtropical regions. In countries such as Brazil, where arboviruses already represent a significant public health burden, climate change has intensified existing vulnerabilities and imposed new demands on health systems, particularly regarding disease surveillance, prevention, and healthcare delivery. This academic essay critically examines the relationship between climate change and the growing burden of arboviral diseases, emphasizing the challenges faced by epidemiological surveillance systems and Primary Health Care (PHC). The discussion explores the ecological mechanisms linking climate variability to vector proliferation, the organizational implications for health services, the importance of integrated surveillance, and the strategic role of PHC in prevention, early diagnosis, community engagement, and intersectoral action. The essay argues that strengthening climate-sensitive health governance and resilient primary healthcare systems is essential for reducing the health impacts of climate-driven arboviral transmission and protecting vulnerable populations.

Keywords: climate change; arboviruses; dengue; epidemiological surveillance; primary health care; public health.

Methodology

This study adopts the format of an academic essay based on critical reflection and theoretical analysis of the scientific literature concerning climate change, arboviral diseases, epidemiological surveillance, and Primary Health Care. Rather than presenting original empirical findings, the manuscript synthesizes current scientific evidence, institutional reports, and public health policy discussions to examine how environmental transformations influence the epidemiology of vector-borne diseases and the organization of healthcare systems. The analytical approach integrates perspectives from epidemiology, environmental health, climate science, health policy, and public health governance.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

The discussion is organized into three thematic sections addressing the ecological relationship between climate change and arboviruses, the challenges faced by epidemiological surveillance systems, and the strategic contribution of Primary Health Care to prevention, preparedness, and community resilience. Particular attention is given to the interaction between environmental determinants, social vulnerability, health system capacity, and intersectoral governance, recognizing that climate-sensitive diseases require coordinated responses extending beyond the traditional healthcare sector.

To enhance readability and maintain a continuous argumentative flow, in-text citations are intentionally omitted throughout the manuscript. All concepts and discussions are grounded in scientific evidence and institutional publications, whose complete bibliographic information is presented in the reference list at the end of the article according to the American Psychological Association (APA, 7th edition). This format prioritizes textual fluency while preserving academic rigor and transparency regarding the sources supporting the analysis.

Introduction

Climate change has become one of the defining global challenges affecting human health, transforming patterns of disease transmission, environmental exposure, and healthcare demand across diverse geographic regions. Rising global temperatures, altered precipitation regimes, more frequent extreme weather events, deforestation, biodiversity loss, and accelerated urbanization have substantially modified the ecological conditions that determine the distribution and behavior of disease vectors. These environmental transformations increasingly influence the occurrence of infectious diseases, particularly those transmitted by mosquitoes, making climate change an essential consideration for contemporary public health planning and disease prevention strategies.

Among climate-sensitive infectious diseases, arboviral illnesses have received growing international attention due to their rapid geographic expansion and increasing epidemiological importance. Diseases such as dengue, chikungunya, Zika, and yellow fever are transmitted primarily by mosquitoes of the *Aedes* genus, whose life cycle, reproductive capacity, survival, and vector competence are directly affected by climatic variables including temperature, rainfall, humidity, and seasonal variation. As these environmental conditions become increasingly favorable in many regions, arboviruses have expanded into areas previously considered unsuitable for sustained transmission, creating new public health challenges for both endemic and non-endemic countries.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Brazil represents one of the countries most affected by this evolving epidemiological scenario. Its tropical climate, continental dimensions, intense urbanization, socioeconomic inequalities, environmental degradation, and widespread distribution of *Aedes aegypti* create highly favorable conditions for recurrent arboviral epidemics. Climate variability has further intensified these challenges by extending transmission seasons, increasing mosquito abundance, and complicating disease surveillance efforts. Simultaneously, Primary Health Care has assumed an increasingly strategic role in early case detection, community education, vector control support, risk communication, and healthcare coordination during outbreaks, highlighting its importance within climate-resilient health systems.

Against this background, this essay addresses the following research question: How does climate change influence the epidemiology of arboviral diseases, and what challenges and opportunities does this transformation create for epidemiological surveillance and Primary Health Care in strengthening climate-resilient public health systems?

Development

1. Climate Change as a Driver of Arboviral Disease Expansion

Climate change has fundamentally altered the ecological conditions that regulate the transmission of vector-borne diseases across the globe. Increasing average temperatures, changing precipitation patterns, prolonged droughts followed by intense rainfall, rising humidity, and the growing frequency of extreme weather events have modified the habitats and reproductive dynamics of numerous mosquito species. These environmental changes have created increasingly favorable conditions for arbovirus transmission, allowing vectors to colonize new territories and sustain longer periods of seasonal activity. Consequently, climate change has become one of the principal environmental determinants influencing the contemporary epidemiology of arboviral diseases.

Arboviruses are transmitted primarily through hematophagous arthropods, especially mosquitoes belonging to the *Aedes* genus. Among these, *Aedes aegypti* is responsible for the transmission of dengue, Zika, chikungunya, and urban yellow fever, while *Aedes albopictus* has emerged as an increasingly important secondary vector in several regions. The biological characteristics of these mosquitoes make them highly responsive to environmental conditions, particularly temperature and moisture, which directly influence their reproductive cycles, survival rates, feeding behavior, and viral replication capacity.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Temperature exerts one of the strongest influences on mosquito biology. Warmer environmental conditions accelerate larval development, shorten the interval between blood meals, increase reproductive activity, and reduce the time required for viral replication within the mosquito. As a result, infected mosquitoes become capable of transmitting arboviruses more rapidly, increasing the efficiency of disease transmission. However, extremely high temperatures may also reduce mosquito survival, illustrating the complex and nonlinear relationship between climate variables and vector ecology.

Rainfall patterns similarly affect mosquito populations through their influence on breeding site availability. Periods of moderate rainfall often increase the accumulation of stagnant water in artificial containers, discarded waste, construction sites, and urban infrastructure, creating ideal environments for mosquito oviposition. Conversely, prolonged droughts may also contribute to transmission by encouraging water storage practices in households, generating numerous artificial breeding sites where vector populations can proliferate. Thus, both excessive rainfall and water scarcity may indirectly favor arboviral transmission under different environmental circumstances.

Humidity represents another important climatic determinant affecting vector survival and dispersal. High humidity generally prolongs mosquito lifespan by reducing dehydration, thereby increasing the probability that infected mosquitoes survive long enough to transmit viruses to susceptible human hosts. Because arboviral transmission depends on the completion of viral incubation within the mosquito, longer vector survival substantially enhances epidemic potential. Consequently, climatic conditions combining elevated temperatures with high humidity often produce particularly favorable environments for arbovirus circulation.

Climate change has also expanded the geographic distribution of mosquito vectors. Regions previously characterized by cooler temperatures or shorter transmission seasons are increasingly becoming environmentally suitable for *Aedes* establishment. Mountainous areas, subtropical regions, and temperate zones have reported the emergence of vector populations that were previously absent or unable to sustain long-term transmission. This geographic expansion exposes immunologically naïve populations to arboviruses, increasing the risk of outbreaks in areas with limited surveillance capacity and little prior experience managing these diseases.

Urbanization interacts closely with climate change in shaping arboviral epidemiology. Rapid urban growth frequently occurs without adequate sanitation infrastructure, waste management, drainage systems, or reliable water supply, creating environments highly conducive to mosquito breeding. Dense human

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

populations facilitate mosquito-human contact while increasing opportunities for viral transmission. Climate-related environmental changes therefore often amplify existing urban vulnerabilities rather than acting as isolated determinants of disease emergence.

Deforestation and ecosystem degradation further contribute to changing transmission dynamics. The destruction of natural habitats alters ecological relationships among wildlife, vectors, and human populations, increasing opportunities for pathogen spillover and modifying mosquito distribution patterns. Environmental disturbances may also reduce biodiversity that naturally regulates vector populations, allowing disease-transmitting mosquitoes to become more abundant in disturbed ecosystems. Consequently, environmental degradation and climate change frequently operate through interconnected ecological mechanisms influencing disease emergence.

Globalization and human mobility intensify the effects of climate change by facilitating the rapid dissemination of arboviruses across international borders. International travel, migration, trade, and urban connectivity allow infected individuals and mosquito vectors to introduce viruses into environmentally suitable regions where local transmission can subsequently become established. Climate change therefore interacts with globalization to accelerate the geographic spread of arboviral diseases, transforming local outbreaks into regional or international public health concerns.

The epidemiology of dengue illustrates the profound influence of climatic variability on disease dynamics. Seasonal increases in temperature and rainfall frequently precede dengue epidemics by creating favorable conditions for mosquito population growth. In many endemic countries, unusually warm or wet seasons are associated with higher case numbers, prolonged transmission periods, and greater epidemic intensity. Climate forecasting has therefore become increasingly valuable for anticipating periods of elevated transmission risk and supporting early public health preparedness.

Zika virus transmission similarly demonstrated the complex interaction between environmental, biological, and social determinants. The explosive epidemic observed throughout the Americas during 2015 and 2016 highlighted how favorable climatic conditions, widespread vector distribution, susceptible populations, and intense human mobility could combine to facilitate rapid continental dissemination. The association between Zika infection during pregnancy and congenital abnormalities further emphasized the far-reaching consequences of climate-sensitive vector-borne diseases beyond immediate infectious morbidity.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Chikungunya has likewise expanded substantially in recent decades, benefiting from environmental conditions favorable to both *Aedes aegypti* and *Aedes albopictus*. Unlike dengue, chikungunya frequently produces debilitating chronic musculoskeletal symptoms that may persist for months or even years, generating prolonged healthcare demand and socioeconomic consequences. Climate-driven increases in transmission therefore carry implications extending beyond acute epidemic response, requiring sustained clinical follow-up and rehabilitation services within health systems.

The health impacts of climate-sensitive arboviruses are not distributed equally across populations. Social vulnerability strongly influences exposure to mosquito breeding sites, housing quality, sanitation access, waste disposal, healthcare utilization, and the capacity to implement preventive measures. Low-income communities frequently experience greater environmental exposure while simultaneously facing barriers to healthcare access, illustrating how climate change interacts with existing social inequalities to produce disproportionate health burdens among disadvantaged populations.

The increasing frequency of climate extremes has further complicated arboviral control. Floods, hurricanes, prolonged droughts, heatwaves, and other environmental disasters may disrupt vector control activities, damage healthcare infrastructure, interrupt surveillance systems, and displace vulnerable populations into conditions favoring disease transmission. Public health preparedness must therefore increasingly incorporate climate adaptation strategies capable of maintaining essential surveillance and healthcare services during environmental emergencies.

Ultimately, climate change should not be viewed merely as an environmental issue but as a major determinant of infectious disease epidemiology and health system resilience. The expansion of arboviral diseases reflects the interaction of climatic variability with urbanization, environmental degradation, globalization, socioeconomic inequalities, and public health capacity. Understanding these interconnected drivers is essential for developing effective prevention strategies and for strengthening surveillance systems and Primary Health Care in an era of accelerating environmental change.

2. Challenges for Epidemiological Surveillance in the Context of Climate Change

Epidemiological surveillance occupies a central position in the prevention and control of arboviral diseases, particularly in the context of accelerating climate

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

change. Surveillance systems are responsible for detecting outbreaks, monitoring disease trends, identifying environmental risk factors, guiding public health interventions, and supporting evidence-based decision-making. However, climate variability has introduced new levels of uncertainty into disease transmission patterns, making surveillance increasingly complex. Traditional surveillance models based primarily on historical epidemiological data are becoming less effective as environmental conditions rapidly evolve and transmission dynamics become more unpredictable.

One of the principal challenges involves the changing seasonality of arboviral diseases. Historically, outbreaks of dengue and other mosquito-borne infections followed relatively predictable seasonal patterns associated with rainy periods and higher temperatures. Climate change, however, has altered these temporal cycles by extending transmission seasons, reducing the duration of low-risk periods, and increasing the occurrence of off-season outbreaks. Consequently, surveillance systems must increasingly operate as continuous, year-round monitoring mechanisms rather than concentrating activities during traditionally recognized epidemic periods.

The geographic expansion of vector populations presents another major surveillance challenge. Regions previously considered unsuitable for sustained *Aedes* mosquito transmission are increasingly reporting vector establishment and locally acquired infections. This expansion requires surveillance authorities to monitor territories that historically lacked specialized entomological expertise, laboratory infrastructure, and trained personnel. As transmission zones become more widespread, surveillance systems must adapt their territorial coverage while strengthening coordination across municipalities, states, and national public health institutions.

Climate change has also increased the frequency of simultaneous outbreaks affecting multiple geographic regions. Rather than isolated epidemics occurring in specific municipalities, public health authorities increasingly confront synchronized transmission across extensive territorial areas. Such widespread epidemics place considerable pressure on surveillance capacity by increasing notification volumes, laboratory demand, epidemiological investigations, and the need for rapid dissemination of information to healthcare providers and decision-makers.

Entomological surveillance has become progressively more important within climate-sensitive disease monitoring. Understanding mosquito density, breeding site distribution, insecticide resistance, species composition, and seasonal population dynamics provides essential information for anticipating epidemic risk. Nevertheless, maintaining comprehensive entomological surveillance requires

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

substantial financial investment, specialized technical expertise, laboratory support, and continuous field operations. Resource limitations frequently restrict the capacity of health systems to conduct systematic vector monitoring at the scale required by expanding transmission areas.

Laboratory surveillance faces similar pressures. Confirming arboviral infections often requires molecular or serological testing capable of distinguishing among viruses with overlapping clinical presentations. During major outbreaks, laboratory services may become overwhelmed by the large volume of suspected cases, delaying diagnosis and reducing the timeliness of epidemiological reporting. Climate-driven increases in arboviral incidence therefore require sustained investment in laboratory capacity, diagnostic technologies, and workforce development to ensure rapid and reliable disease confirmation.

The clinical similarities among dengue, Zika, chikungunya, and other febrile illnesses further complicate surveillance activities. Patients frequently present with nonspecific symptoms such as fever, headache, rash, muscle pain, and fatigue, making clinical differentiation difficult, particularly during the early stages of illness. Co-circulation of multiple arboviruses may therefore result in underreporting, diagnostic uncertainty, or delayed recognition of emerging outbreaks. Accurate surveillance increasingly depends upon integrating clinical assessment with laboratory confirmation and epidemiological investigation.

Data quality represents another critical determinant of surveillance effectiveness. Timely notification, complete reporting, standardized case definitions, and reliable health information systems are fundamental for accurately describing transmission dynamics. Incomplete records, delayed reporting, inconsistent data collection practices, and fragmented information systems may compromise situational awareness, reducing the ability of public health authorities to detect outbreaks promptly and implement appropriate interventions. Strengthening data governance has therefore become an essential component of climate-resilient surveillance systems.

The integration of environmental and epidemiological information has emerged as a strategic priority. Climate variables such as temperature, rainfall, humidity, vegetation indices, and extreme weather events provide valuable indicators capable of anticipating periods of increased transmission risk. Incorporating meteorological and environmental data into disease surveillance allows health authorities to move from reactive outbreak response toward more proactive risk prediction and preparedness. Such integration requires close collaboration between public health agencies, meteorological institutions, environmental authorities, and academic researchers.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Technological innovation offers significant opportunities for strengthening surveillance capacity. Geographic information systems, remote sensing, satellite imagery, artificial intelligence, machine learning, predictive modeling, and digital reporting platforms increasingly support the analysis of large and complex datasets. These technologies may improve the identification of high-risk areas, facilitate real-time monitoring, and generate early warning systems capable of anticipating outbreaks before widespread transmission occurs. Nevertheless, successful implementation depends upon adequate digital infrastructure, technical expertise, and ethical governance of health data.

Community participation also represents a valuable component of surveillance. Citizens frequently serve as the first observers of increasing mosquito populations, unusual disease occurrence, or environmental conditions favorable to vector proliferation. Community reporting mechanisms, health education initiatives, and participatory surveillance approaches can complement formal epidemiological systems by improving the early identification of local risks. Building public trust and encouraging active community engagement strengthen surveillance while promoting shared responsibility for disease prevention.

Climate change has further highlighted the importance of adopting the One Health perspective within surveillance systems. Human health, animal health, and environmental conditions are deeply interconnected, particularly regarding vector-borne diseases whose transmission depends upon ecological processes. Surveillance strategies that integrate environmental monitoring, entomological data, veterinary information, and human disease reporting provide a more comprehensive understanding of transmission dynamics than isolated sectoral approaches. Interdisciplinary collaboration therefore becomes increasingly essential for addressing climate-sensitive health threats.

International cooperation has gained growing relevance as arboviruses expand across national borders. Mosquito vectors and infectious diseases do not recognize political boundaries, making regional collaboration indispensable for surveillance, laboratory networking, information exchange, and coordinated outbreak response. International organizations increasingly promote standardized surveillance methodologies, shared reporting systems, and collaborative research capable of improving preparedness for emerging vector-borne diseases influenced by global climate change.

Financial sustainability remains an underlying challenge affecting every dimension of epidemiological surveillance. Maintaining laboratory networks, information systems, vector monitoring, professional training, technological innovation, and emergency preparedness requires continuous public investment.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Surveillance systems often receive increased attention during epidemics but may experience reduced funding once outbreaks decline. Climate change, however, demands permanent surveillance capacity because environmental conditions increasingly support year-round transmission and recurrent epidemic cycles.

Epidemiological surveillance must evolve from a predominantly reactive system into a proactive, integrated, climate-informed public health function. Future surveillance models should combine epidemiological intelligence, environmental monitoring, technological innovation, community participation, and intersectoral collaboration to anticipate emerging risks before widespread outbreaks occur. Such transformation is essential for strengthening health system resilience and protecting populations from the growing public health consequences of climate-driven arboviral diseases.

3. The Strategic Role of Primary Health Care in Climate-Resilient Arboviral Disease Control

Primary Health Care (PHC) occupies a strategic position in addressing the growing burden of arboviral diseases associated with climate change. As the first level of contact between individuals and the healthcare system, PHC provides comprehensive, continuous, and community-oriented care capable of integrating prevention, health promotion, surveillance, diagnosis, treatment, and long-term follow-up. Within climate-sensitive health systems, strengthening Primary Health Care is essential not only for improving clinical outcomes but also for increasing community resilience and reducing the social impacts of recurrent arboviral epidemics.

One of the principal contributions of PHC lies in its capacity for early case identification. Patients experiencing the initial symptoms of dengue, chikungunya, or Zika commonly seek care at primary healthcare facilities before requiring specialized treatment. Early clinical recognition allows healthcare professionals to initiate appropriate management, identify warning signs, monitor disease progression, and refer severe cases when necessary. Timely diagnosis also contributes to more accurate epidemiological surveillance by improving disease notification and facilitating rapid public health responses.

Health promotion constitutes another fundamental responsibility of Primary Health Care. Community education regarding mosquito breeding sites, household sanitation, personal protective measures, recognition of early symptoms, and appropriate healthcare-seeking behavior significantly contributes to reducing disease transmission. Educational initiatives are particularly effective when adapted

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

to local cultural contexts and implemented through continuous dialogue with communities rather than isolated information campaigns conducted only during epidemic periods.

Community Health Workers play an especially important role in climate-sensitive arboviral control. Through regular household visits, these professionals establish close relationships with families, identify environmental risk factors, encourage preventive behaviors, monitor vulnerable individuals, and strengthen communication between healthcare services and local communities. Their territorial knowledge enables the identification of neighborhoods requiring intensified vector control activities, thereby improving the responsiveness of public health interventions.

The Family Health Strategy further enhances the contribution of Primary Health Care by organizing multidisciplinary teams responsible for defined geographic territories. This territorial approach facilitates continuous monitoring of environmental conditions, demographic changes, and epidemiological patterns affecting specific communities. Because healthcare professionals maintain long-term relationships with local populations, they are better positioned to recognize emerging outbreaks, coordinate preventive activities, and mobilize community participation in vector control efforts.

Climate change increasingly requires PHC to adopt a proactive rather than reactive approach. Instead of responding only after outbreaks have become established, primary healthcare teams can identify climatic and environmental conditions associated with increased transmission risk and implement preventive actions before widespread disease occurs. Seasonal planning, risk mapping, community mobilization, and intensified surveillance during high-risk periods contribute to greater preparedness and more effective epidemic prevention.

Integrated management of chronic diseases also becomes increasingly important during arboviral epidemics. Older adults, pregnant women, children, and individuals living with chronic illnesses frequently experience greater vulnerability to severe complications. Primary healthcare professionals are uniquely positioned to monitor these high-risk populations, ensure continuity of essential healthcare services, and provide individualized guidance during periods of increased arboviral transmission. Maintaining routine healthcare while responding to outbreaks prevents indirect health consequences associated with interruptions in chronic disease management.

Mental health support represents another frequently overlooked dimension of Primary Health Care during arboviral epidemics. Large outbreaks often generate fear, anxiety, uncertainty, and psychological distress within affected communities, particularly among pregnant women concerned about congenital infections or

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

individuals experiencing prolonged complications following chikungunya infection. Primary healthcare services can provide psychosocial support, risk communication, and community-based mental health interventions that strengthen resilience during public health emergencies.

Intersectoral collaboration significantly enhances the effectiveness of Primary Health Care in addressing climate-sensitive diseases. Successful arboviral prevention depends not only on healthcare services but also on sanitation, waste management, urban planning, environmental protection, education, housing policy, and community development. Primary healthcare teams frequently serve as connectors among these sectors by identifying local environmental problems and facilitating collaborative actions involving governmental agencies, schools, community organizations, and civil society.

Digital health technologies increasingly support the work of Primary Health Care professionals. Electronic health records, telemedicine, mobile health applications, geographic information systems, and digital surveillance platforms improve communication, facilitate case reporting, enhance continuity of care, and strengthen coordination with epidemiological surveillance authorities. When appropriately implemented, these technologies expand the capacity of PHC to monitor arboviral transmission while improving access to healthcare services, particularly in geographically isolated communities.

Professional education and continuous training are indispensable for maintaining effective Primary Health Care responses to climate-sensitive diseases. Healthcare professionals must remain updated regarding evolving clinical guidelines, emerging arboviruses, diagnostic criteria, vector ecology, climate-health interactions, and risk communication strategies. Continuing education programs strengthen clinical competence while improving the ability of multidisciplinary teams to respond to increasingly complex epidemiological scenarios associated with environmental change.

Primary Health Care also contributes to scientific knowledge generation through routine data collection, surveillance activities, and participation in implementation research. Information generated by frontline healthcare services provides valuable insights into disease distribution, healthcare utilization, intervention effectiveness, and community needs. Integrating primary care data with epidemiological and environmental information strengthens evidence-based policymaking while supporting continuous improvement of climate adaptation strategies.

Health equity should remain a central objective of climate-resilient Primary Health Care. Climate change disproportionately affects socially vulnerable populations

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

living in inadequate housing, flood-prone neighborhoods, areas with deficient sanitation, and communities with limited access to healthcare services. Primary healthcare teams play a critical role in identifying these vulnerable groups, prioritizing preventive interventions, and reducing inequalities in healthcare access. Equity-oriented strategies ensure that climate adaptation efforts protect those populations facing the greatest environmental and social risks.

Strengthening institutional resilience has become an essential priority for Primary Health Care. Climate change is expected to increase the frequency of extreme weather events capable of disrupting healthcare infrastructure, transportation systems, electricity supply, and communication networks. Resilient primary healthcare services require emergency preparedness plans, adequate infrastructure, supply chain management, workforce flexibility, and contingency protocols that allow essential services to continue operating during environmental crises and disease outbreaks.

Primary Health Care represents the cornerstone of climate-resilient health systems capable of responding effectively to the expanding burden of arboviral diseases. By integrating prevention, surveillance, community engagement, multidisciplinary care, intersectoral collaboration, digital innovation, and social participation, PHC strengthens both individual health outcomes and collective public health preparedness. As climate change continues to reshape the epidemiology of infectious diseases, investing in robust, equitable, and resilient Primary Health Care will remain one of the most effective strategies for protecting populations and ensuring sustainable public health responses.

Conclusion

Climate change has emerged as one of the most significant environmental determinants influencing the global expansion of arboviral diseases. Rising temperatures, altered precipitation patterns, increasing humidity, ecosystem degradation, rapid urbanization, and extreme weather events have collectively transformed the ecological conditions that favor mosquito proliferation and virus transmission. These environmental changes have extended transmission seasons, expanded the geographic distribution of vectors, and increased the frequency and intensity of outbreaks, particularly in tropical and subtropical countries such as Brazil. In this context, arboviral diseases can no longer be understood solely as infectious diseases requiring vector control but must also be recognized as manifestations of broader climate-related health risks that demand integrated environmental, epidemiological, and healthcare responses. Recent global evidence

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

continues to show that climate variability is contributing to the unprecedented expansion of dengue and other arboviruses while placing increasing pressure on surveillance systems and health services.

Addressing these challenges requires strengthening epidemiological surveillance, investing in climate-informed early warning systems, expanding digital health technologies, and reinforcing Primary Health Care as the cornerstone of climate-resilient health systems. Community engagement, intersectoral governance, environmental management, workforce training, scientific research, and international cooperation should become integral components of national adaptation strategies. Rather than responding only after epidemics emerge, health systems must increasingly anticipate climate-sensitive risks through integrated surveillance, preventive interventions, and evidence-informed public policies. As climate change continues to reshape the epidemiology of vector-borne diseases, resilient Primary Health Care and adaptive public health governance will be indispensable for protecting vulnerable populations, reducing health inequalities, and ensuring sustainable responses to one of the most pressing public health challenges of the twenty-first century.

References

Campbell-Lendrum, D., & Corvalán, C. (2007). Climate change and developing-country cities: Implications for environmental health and equity. *Journal of Urban Health, 84*(Suppl. 1), i109–i117.

Ebi, K. L., & Nealon, J. (2016). Dengue in a changing climate. *Environmental Research, 151*, 115–123.

Intergovernmental Panel on Climate Change. (2023). *Climate change 2023: Synthesis report*. IPCC.

Lowe, R., Barcellos, C., Coelho, C. A. S., Bailey, T. C., Coelho, G. E., Graham, R., & Jupp, T. (2014). Dengue outlook for the World Cup in Brazil: An early warning model framework driven by real-time seasonal climate forecasts. *The Lancet Infectious Diseases, 14*(7), 619–626.

Mordecai, E. A., Ryan, S. J., Caldwell, J. M., Shah, M. M., & LaBeaud, A. D. (2020). Climate change could shift disease burden from malaria to arboviruses in Africa. *The Lancet Planetary Health, 4*(9), e416–e423.

Organização Pan-Americana da Saúde. (2023). *Estratégia para prevenção e controle das arboviroses nas Américas*. OPAS.

CC BY

RSV
ISSN 2674-8584

Received: 21/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/s7ntf066
Pages: 1-16

Ryan, S. J., Carlson, C. J., Mordecai, E. A., & Johnson, L. R. (2019). Global expansion and redistribution of *Aedes*-borne virus transmission risk with climate change. *PLoS Neglected Tropical Diseases, 13*(3), e0007213.

United Nations Environment Programme. (2023). *Adaptation Gap Report 2023*. UNEP.

World Health Organization. (2022). *Global vector control response 2017–2030: Progress report 2022*. World Health Organization.

World Health Organization. (2023). *Climate change and health*. World Health Organization.

World Health Organization. (2023). *Operational framework for building climate resilient and low carbon health systems*. World Health Organization.

World Health Organization. (2024). *Dengue and severe dengue*. World Health Organization.

World Health Organization. (2024). *Global Arbovirus Initiative: Strategic Framework 2022–2030*. World Health Organization.

World Meteorological Organization. (2023). *State of the Global Climate 2023*. WMO.

Xu, Z., Bambrick, H., Yakob, L., Devine, G., Frentiu, F. D., Marina, R., & Hu, W. (2020). The impact of climate change on arboviral disease transmission: A systematic review. *One Health, 11*, 100188.

CC BY