RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Promotion of Physical Activity and Healthy Lifestyle Behaviors as Strategies for Reducing Obesity and Non-Communicable Chronic Diseases

Helen Regina Santos Vitorino
Mestre em Saúde Ambiental e Saúde do Trabalhador
Universidade Federal de Uberlândia, Brasil

Ana Paula Martins de Souza
Mestre em Educação
Universidad Europea del Atlántico

Suzana Maria Teixeira
Técnica de Enfermagem
Universidade Federal de Uberlândia, Brasil

Sirléia Alves de Lima
Técnica de Enfermagem
Universidade Federal de Uberlândia, Brasil

Josina Alves dos Santos
Técnica de Enfermagem
Universidade Federal de Uberlândia, Brasil

Mayara Anna de oliveira Medeiros
Técnica de Enfermagem
Universidade Federal de Uberlândia, Brasil

Fabiane Cristina de Souza Rodovalho
Enfermeira
Universidade Federal de Uberlândia, Brasil

Juliana Cascaes de Aquino Schneider
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

Jocilane Lima de Almeida Vasconcelos
Doutoranda em Saúde Pública
Universidad de Ciencias Empresariales y Sociales, Argentina

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Abstract

Obesity and non-communicable chronic diseases (NCDs) represent some of the greatest public health challenges of contemporary societies. The progressive increase in sedentary behaviors, unhealthy dietary patterns, and social environments that favor excessive caloric intake has contributed to rising rates of obesity, diabetes mellitus, cardiovascular diseases, and other chronic conditions worldwide. Beyond being individual health problems, these conditions are influenced by complex interactions between biological, behavioral, environmental, economic, and social determinants. In this context, promoting physical activity and healthy lifestyle behaviors has become a central strategy for preventing chronic diseases, improving population health, and reducing healthcare system burdens. This academic essay analyzes the role of physical activity promotion and healthy lifestyle behaviors as strategies for reducing obesity and non-communicable chronic diseases. The discussion explores obesity as a public health challenge, the protective effects of regular physical activity and healthy behaviors, and the contribution of Primary Health Care to sustainable health promotion initiatives. The manuscript emphasizes that effective prevention requires integrated approaches combining individual empowerment, supportive environments, public policies, and healthcare interventions capable of addressing the multiple determinants associated with chronic disease development.

Keywords: physical activity; obesity; healthy lifestyle; non-communicable diseases; health promotion; primary health care.

1. Introduction

Obesity has become one of the most important public health challenges of the twenty-first century, affecting populations across different socioeconomic contexts and geographic regions. Once considered mainly a consequence of individual behaviors, obesity is now recognized as a complex chronic condition influenced by biological susceptibility, social environments, economic factors, cultural patterns, and broader structural determinants. Its rapid global expansion has contributed significantly to the increasing prevalence of non-communicable chronic diseases, creating substantial impacts on individual well-being and healthcare systems.

Non-communicable diseases are responsible for a large proportion of global morbidity and mortality, including cardiovascular diseases, diabetes mellitus, chronic respiratory diseases, and several types of cancer. Although genetic factors influence individual vulnerability, many of the determinants associated with these conditions are modifiable, particularly physical inactivity, unhealthy dietary patterns, excessive consumption of ultra-processed foods, tobacco use, and harmful alcohol consumption. Therefore, preventive strategies focused on promoting healthier lifestyles represent essential components of contemporary public health approaches.

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Regular physical activity and healthy behaviors provide significant benefits for preventing and controlling obesity and chronic diseases. Beyond contributing to weight management, physical activity improves cardiovascular function, insulin sensitivity, mental health, functional capacity, and overall quality of life. Similarly, balanced nutrition, adequate sleep, stress management, and avoidance of harmful behaviors contribute to maintaining metabolic health and reducing the risk of chronic disease development. However, individual choices are strongly influenced by social and environmental conditions, making health promotion dependent on supportive policies and accessible opportunities for healthy living.

Considering the increasing burden of obesity and non-communicable chronic diseases worldwide, this essay seeks to answer the following research question: How can the promotion of physical activity and healthy lifestyle behaviors contribute to reducing obesity and non-communicable chronic diseases while strengthening preventive healthcare strategies within Primary Health Care?

2. Methodology

This manuscript is structured as an academic essay based on critical reflection and theoretical analysis of scientific literature addressing obesity, physical activity, healthy behaviors, non-communicable chronic diseases, and health promotion strategies. The objective is not to present original empirical findings but to discuss contemporary concepts, challenges, and opportunities related to lifestyle-based prevention approaches within public health and healthcare systems.

The discussion is organized into three thematic sections. The first section examines obesity and non-communicable diseases as interconnected public health challenges influenced by behavioral, environmental, and social determinants. The second section analyzes the role of physical activity and healthy lifestyle promotion as preventive strategies capable of reducing chronic disease risks. The third section discusses the contribution of Primary Health Care in implementing health promotion actions, supporting behavioral change, and integrating lifestyle interventions into routine healthcare practices.

To preserve argumentative continuity and improve reading fluency, scientific references are not inserted throughout the body of the manuscript. All concepts, theoretical discussions, and evidence presented in this essay are supported by peer-reviewed publications, international reports, and scientific guidelines, which are listed in full in the reference section according to the American Psychological Association (APA, 7th edition) formatting style.

3. Development

3.1 Obesity and Non-Communicable Diseases as a Global Public Health Challenge

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Obesity has become one of the most prominent health challenges worldwide due to its rapid expansion, chronic nature, and association with multiple adverse health outcomes. The increasing prevalence of excessive body weight reflects profound transformations in contemporary societies, including changes in food systems, urban environments, work patterns, transportation habits, and daily routines. Unlike isolated conditions caused by a single factor, obesity emerges from complex interactions between biological predisposition, behavioral practices, social circumstances, and environmental influences that shape individual and collective health.

The global increase in obesity has occurred alongside major epidemiological transitions characterized by longer life expectancy and the growing predominance of non-communicable diseases. As infectious diseases have declined in many regions, chronic conditions have become the leading causes of disability and premature mortality. Obesity contributes significantly to this transition because it increases the risk of several chronic disorders and amplifies the complexity of healthcare needs among affected populations.

Excess body weight is strongly associated with metabolic alterations that contribute to the development of chronic diseases. Increased adipose tissue accumulation promotes inflammatory processes, hormonal changes, insulin resistance, and disturbances in lipid metabolism. These mechanisms create favorable conditions for the emergence of diabetes mellitus, hypertension, cardiovascular diseases, and other long-term health problems. Therefore, obesity should not be understood merely as an aesthetic concern but as a complex chronic condition with significant physiological consequences.

The relationship between obesity and diabetes mellitus represents one of the most important public health concerns associated with unhealthy lifestyles. Excess body weight increases the likelihood of impaired glucose regulation and progressive metabolic dysfunction, contributing to the growing prevalence of type 2 diabetes. As diabetes itself represents a major risk factor for cardiovascular disease, kidney disease, vision impairment, and other complications, preventing obesity becomes a fundamental strategy for reducing the broader burden of chronic illness.

Cardiovascular diseases represent another major consequence of obesity and associated metabolic disorders. Excessive adiposity contributes to increased blood pressure, endothelial dysfunction, chronic inflammation, and unfavorable lipid profiles, creating conditions that accelerate vascular damage. The coexistence of obesity with other risk factors such as physical inactivity, poor nutrition, and smoking further increases cardiovascular risk, demonstrating the importance of integrated prevention strategies rather than isolated interventions targeting individual conditions.

The impact of obesity extends beyond physical health outcomes. Individuals living with obesity frequently experience psychological distress, reduced quality of life, social stigma, discrimination, and barriers to healthcare access. Weight-related stigma may discourage individuals from seeking medical support, reduce engagement with preventive services, and negatively affect mental health. Addressing obesity therefore requires

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

approaches that promote dignity, inclusion, and supportive environments rather than focusing exclusively on individual responsibility.

Social determinants play a central role in shaping patterns of obesity and chronic disease. Income level, education, employment conditions, neighborhood characteristics, food availability, and access to recreational spaces influence the opportunities individuals have to maintain healthy behaviors. Populations experiencing socioeconomic disadvantage often face greater exposure to environments that encourage unhealthy eating and sedentary lifestyles while simultaneously experiencing reduced access to preventive healthcare and health promotion resources.

The modern food environment has significantly influenced patterns of nutrition and obesity development. Increased availability and affordability of ultra-processed foods, aggressive marketing strategies, larger portion sizes, and changing dietary habits have contributed to excessive caloric intake in many populations. These transformations demonstrate that food choices are not determined solely by personal preferences but are also shaped by economic, cultural, and commercial forces operating within broader social systems.

Urbanization has also transformed the way people move, work, and interact with their environments. Many urban areas have been designed around motorized transportation rather than active mobility, reducing opportunities for walking and cycling. Limited access to safe public spaces, parks, and recreational facilities can discourage physical activity, particularly among vulnerable populations. Creating healthier urban environments is therefore an essential component of obesity prevention strategies.

Sedentary behavior has become increasingly common due to technological advances, changes in occupational structures, and increased screen time. Many individuals spend prolonged periods sitting during work, transportation, leisure activities, and daily routines. This reduction in movement has important consequences for metabolic health, cardiovascular function, and physical capacity, even among individuals who occasionally engage in structured exercise. Addressing sedentary lifestyles requires interventions that promote movement throughout the entire day.

Children and adolescents represent a particularly important population for obesity prevention. Early-life exposure to unhealthy dietary patterns, insufficient physical activity, excessive screen time, and obesogenic environments increases the likelihood of obesity persistence into adulthood. Childhood obesity is associated with immediate health consequences and increases the risk of developing chronic diseases later in life. Preventive actions targeting younger populations therefore provide long-term benefits for individuals and societies.

Population ageing introduces additional challenges related to obesity and chronic disease management. Older adults may experience reduced mobility, loss of muscle mass, functional limitations, and multiple chronic conditions that complicate weight management and physical activity participation. Prevention strategies must therefore consider different

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

life stages and adapt recommendations according to individual capacities, health conditions, and social circumstances.

The economic consequences of obesity and chronic diseases place considerable pressure on healthcare systems. Costs arise from medical consultations, medications, hospitalizations, treatment of complications, disability, and productivity losses. As chronic diseases require long-term management, prevention strategies capable of reducing disease incidence and progression offer significant benefits for healthcare sustainability and resource allocation.

Effective responses to obesity require moving beyond approaches focused exclusively on individual behavior change. Although personal choices influence health outcomes, these choices occur within social and environmental contexts that can either facilitate or restrict healthy behaviors. Sustainable prevention requires coordinated actions involving healthcare systems, educational institutions, urban planning, food policies, workplaces, communities, and governmental organizations.

From a public health perspective, obesity and non-communicable diseases represent interconnected challenges requiring comprehensive and long-term strategies. Addressing these conditions demands interventions capable of modifying environments, strengthening health promotion, improving access to healthy choices, and empowering individuals to adopt sustainable behaviors. Recognizing obesity as a complex social and health issue is essential for developing effective policies that reduce chronic disease burden and promote healthier populations.

3.2 Physical Activity and Healthy Lifestyle Promotion as Preventive Strategies

Physical activity represents one of the most effective strategies for preventing and managing obesity and non-communicable chronic diseases. Regular movement contributes to improvements in cardiovascular health, metabolic regulation, muscular strength, functional capacity, and psychological well-being. Unlike many preventive interventions that require complex technologies or specialized resources, physical activity can generate substantial health benefits through accessible and adaptable practices incorporated into daily routines. For this reason, promoting active lifestyles has become a central component of global strategies aimed at reducing the burden of chronic diseases.

The benefits of physical activity extend far beyond body weight control. Although increased energy expenditure contributes to weight management, regular movement also produces independent physiological effects that improve insulin sensitivity, blood pressure regulation, lipid metabolism, inflammatory processes, and cardiovascular function. These effects demonstrate that physical activity should not be considered exclusively as a tool for weight reduction but as a broader intervention capable of improving overall health regardless of changes in body composition.

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Different forms of physical activity provide important health benefits throughout the life course. Aerobic activities, resistance exercises, flexibility practices, and activities that improve balance contribute to maintaining physical capacity and preventing functional decline. The most effective approach is often one that considers individual preferences, health conditions, cultural contexts, and available resources, increasing the likelihood that healthy behaviors will be maintained over time.

The relationship between physical activity and diabetes prevention represents a major area of public health interest. Regular movement improves glucose utilization by skeletal muscles, enhances insulin sensitivity, and contributes to better metabolic control among individuals at risk or already living with diabetes. These effects are particularly relevant considering the global increase in type 2 diabetes associated with population ageing, obesity, and sedentary lifestyles.

Cardiovascular health also benefits substantially from regular physical activity. Active lifestyles contribute to improved vascular function, reduced blood pressure, better lipid profiles, and enhanced cardiac capacity. These protective effects reduce the likelihood of cardiovascular events and support healthier ageing. Encouraging physical activity among individuals with cardiovascular risk factors should therefore be considered an essential component of chronic disease prevention and management.

Healthy eating patterns represent another fundamental component of chronic disease prevention. Adequate nutrition contributes to maintaining healthy body weight, regulating metabolism, supporting immune function, and preventing nutritional deficiencies. However, promoting healthy diets requires consideration of broader factors such as food accessibility, cultural preferences, economic conditions, and the availability of nutritious options within communities.

The quality of dietary patterns has become increasingly important in the context of modern food systems. The widespread consumption of ultra-processed foods, excessive intake of added sugars, high sodium consumption, and insufficient intake of fruits, vegetables, and whole foods are associated with increased risks of obesity and chronic diseases. Improving population nutrition therefore requires both individual education and structural interventions capable of transforming food environments.

Sleep quality and stress management have also gained recognition as important components of healthy lifestyles. Insufficient sleep duration, irregular sleep patterns, and chronic psychological stress may influence appetite regulation, hormonal balance, emotional well-being, and metabolic health. Lifestyle promotion strategies that include adequate sleep and stress reduction provide a more comprehensive approach than interventions focused exclusively on physical activity and diet.

The reduction of sedentary behavior represents an additional priority in health promotion. Even individuals who engage in regular exercise may experience negative health effects when spending excessive amounts of time sitting throughout the day. Encouraging frequent movement breaks, active transportation, workplace activity, and

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

reduced prolonged screen time contributes to healthier daily routines and complements structured exercise programs.

Behavioral change is a complex process influenced by motivation, knowledge, social support, habits, and environmental conditions. Health promotion interventions are more effective when they recognize that adopting healthy behaviors requires sustained support rather than temporary encouragement. Healthcare professionals play an important role in helping individuals establish realistic goals, overcome barriers, and develop strategies compatible with their daily circumstances.

The social environment strongly influences the ability of individuals to maintain healthy behaviors. Family support, community networks, workplace conditions, and cultural norms can either facilitate or hinder participation in physical activity and healthy eating practices. Interventions that involve social groups and community participation often produce more sustainable results because they transform health promotion from an individual responsibility into a collective process.

Schools represent strategic environments for promoting healthy lifestyles from early childhood. Educational institutions can contribute by incorporating physical activity opportunities, nutrition education, healthier food options, and initiatives that encourage positive relationships with movement and body image. Early interventions help establish lifelong habits while preventing the development of obesity and chronic diseases during later stages of life.

Workplaces also offer important opportunities for health promotion among adults. Occupational environments influence daily movement patterns, stress levels, eating habits, and opportunities for preventive care. Workplace interventions that encourage active breaks, healthier food choices, ergonomic improvements, and wellness initiatives can positively influence employee health while reducing productivity losses associated with chronic diseases.

Community-based approaches are essential for reducing inequalities in access to healthy lifestyles. Individuals living in disadvantaged communities may face barriers such as unsafe environments, limited recreational spaces, financial constraints, and restricted access to nutritious foods. Public policies that create supportive environments, improve infrastructure, and expand opportunities for physical activity contribute to reducing health disparities and promoting more equitable outcomes.

Technology has created new possibilities for encouraging physical activity and healthy behaviors. Digital applications, wearable devices, online educational resources, and remote monitoring tools can support self-management, provide feedback, and increase engagement with preventive strategies. Nevertheless, technological approaches should complement rather than replace human interaction and community-based interventions, particularly among populations experiencing digital inequalities.

Taken together, physical activity and healthy lifestyle promotion represent powerful strategies for reducing obesity and non-communicable chronic diseases. Their

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

effectiveness depends not only on individual adoption of healthier behaviors but also on the creation of social and environmental conditions that make these choices accessible and sustainable. Long-term improvements in population health require coordinated actions that combine education, supportive policies, community participation, and healthcare interventions capable of promoting healthier ways of living across the life course.

3.3 The Role of Primary Health Care in Promoting Healthy Behaviors and Reducing Chronic Disease Burden

Primary Health Care occupies a central position in strategies aimed at promoting healthy lifestyles and reducing the burden of obesity and non-communicable chronic diseases. As the most accessible level of healthcare, primary care services maintain continuous relationships with individuals, families, and communities, creating opportunities for prevention, health education, early intervention, and long-term support. Unlike healthcare models focused primarily on treating established diseases, Primary Health Care has the potential to address risk factors before they develop into complex clinical conditions.

The preventive nature of Primary Health Care makes it a strategic setting for promoting physical activity and healthy behaviors. Routine consultations provide opportunities to assess lifestyle patterns, identify risk factors, establish individualized goals, and support gradual behavioral changes. Integrating health promotion into regular care ensures that prevention becomes part of everyday clinical practice rather than an occasional activity implemented only during public health campaigns.

Healthcare professionals working in Primary Health Care play an important role in encouraging individuals to adopt healthier lifestyles. Physicians, nurses, nutritionists, community health workers, psychologists, and other professionals contribute different perspectives to understanding and addressing the multiple factors associated with obesity and chronic diseases. This multidisciplinary approach recognizes that lifestyle change involves biological, psychological, social, and environmental dimensions that cannot be effectively addressed through isolated interventions.

Health counseling represents one of the most common strategies used within primary care to promote behavioral change. Effective counseling involves more than providing general recommendations; it requires understanding individual circumstances, identifying barriers, negotiating realistic objectives, and supporting patients throughout the process of change. Personalized approaches increase the likelihood that individuals will adopt sustainable behaviors according to their abilities, preferences, and social realities.

Community Health Workers provide a valuable connection between healthcare services and local populations. Through their proximity to households and communities, they can identify social and environmental factors influencing health behaviors, encourage participation in preventive activities, and strengthen communication between healthcare

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

teams and residents. Their territorial knowledge allows interventions to be adapted according to specific community needs and cultural contexts.

Primary Health Care also contributes to obesity prevention through the early identification of individuals at increased risk. Monitoring body weight, nutritional status, physical activity patterns, blood pressure, and metabolic indicators enables healthcare professionals to recognize risk trajectories before severe complications develop. Early intervention allows preventive actions to be initiated when behavioral changes may have greater impact and when individuals may still maintain preserved health conditions.

The management of obesity within Primary Health Care requires approaches that avoid stigma and emphasize comprehensive health improvement rather than focusing exclusively on body weight. Weight-related stigma may negatively affect psychological well-being, reduce trust in healthcare professionals, and discourage individuals from seeking assistance. Supportive communication that recognizes obesity as a complex chronic condition contributes to more respectful and effective care.

Health promotion activities developed through Primary Health Care should extend beyond individual consultations. Group activities, educational workshops, community exercise programs, cooking initiatives, and partnerships with local organizations can create supportive environments for healthy behaviors. These collective strategies strengthen social connections and provide opportunities for individuals to share experiences, challenges, and strategies for maintaining healthier lifestyles.

Intersectoral collaboration is essential for addressing the broader determinants associated with obesity and chronic diseases. Healthcare services alone cannot modify all factors influencing lifestyle behaviors. Cooperation with schools, workplaces, urban planning sectors, social assistance programs, and community organizations is necessary to create environments that facilitate physical activity, improve food access, and promote healthier daily routines.

The physical environment surrounding communities has a significant influence on health behaviors. Primary Health Care teams can contribute to identifying local barriers such as lack of recreational spaces, unsafe streets, limited access to healthy foods, or transportation difficulties. By communicating these challenges to other sectors and participating in local health planning, primary care services can contribute to broader community-level interventions.

Digital health resources have expanded the possibilities for lifestyle promotion within Primary Health Care. Telehealth services, mobile applications, digital educational materials, and remote monitoring tools can complement face-to-face care and increase access to health promotion strategies. These resources are particularly valuable for individuals who experience geographic, occupational, or mobility-related barriers to accessing traditional healthcare services.

Continuous professional training is necessary to ensure that healthcare teams are prepared to address obesity and lifestyle-related chronic diseases effectively. Professionals

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

require updated knowledge regarding behavior change strategies, nutritional counseling, physical activity recommendations, motivational approaches, and the social determinants influencing health behaviors. Investment in workforce development strengthens the capacity of Primary Health Care to deliver comprehensive preventive care.

Healthcare systems should also incorporate evaluation mechanisms to monitor the effectiveness of health promotion initiatives. Measuring indicators related to physical activity levels, nutritional behaviors, obesity prevalence, metabolic outcomes, and patient engagement allows services to identify successful approaches and improve interventions over time. Continuous assessment transforms health promotion from isolated activities into organized and evidence-informed strategies.

Equity must remain a fundamental principle in Primary Health Care approaches to obesity prevention and chronic disease reduction. Populations facing socioeconomic disadvantage frequently encounter greater exposure to unhealthy environments and fewer opportunities to engage in preventive behaviors. Primary care services have an important responsibility in identifying these inequalities and developing interventions capable of expanding access to health-promoting resources.

The integration of lifestyle promotion into chronic disease management also contributes to more efficient healthcare systems. Preventing obesity-related complications reduces the need for expensive treatments, hospitalizations, and long-term disease management. By focusing on prevention and early intervention, Primary Health Care supports both improved population health outcomes and greater sustainability of healthcare resources.

Considering these aspects, Primary Health Care should be recognized as a key platform for transforming health promotion from a theoretical principle into practical action. Through continuous care, multidisciplinary collaboration, community engagement, and preventive interventions, primary care services can significantly influence patterns of physical activity, nutrition, and chronic disease risk. Strengthening this level of healthcare is therefore essential for creating healthier communities and reducing the long-term burden of obesity and non-communicable chronic diseases.

4. Conclusion

Obesity and non-communicable chronic diseases represent complex and interconnected public health challenges influenced by individual behaviors, social determinants, environmental conditions, and broader structural factors. The increasing prevalence of sedentary lifestyles, unhealthy dietary patterns, and obesogenic environments has contributed to a substantial rise in chronic disease burden worldwide, affecting healthcare systems, economies, and population well-being. In this context, promoting physical activity and healthy lifestyle behaviors represents a fundamental strategy for preventing disease, improving quality of life, and reducing avoidable health complications throughout the life course.

CC BY

RSV
ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Primary Health Care plays a decisive role in translating health promotion principles into effective and sustainable interventions. Through continuous patient relationships, multidisciplinary teamwork, community engagement, and integration of preventive actions into routine care, primary healthcare services can support individuals in adopting healthier behaviors while addressing the social and environmental factors that influence health choices. Reducing obesity and chronic diseases requires coordinated efforts involving healthcare systems, public policies, communities, educational institutions, workplaces, and other sectors of society. Building environments that facilitate healthy living is therefore essential for achieving long-term improvements in population health.

References

Bennett, J. E., Stevens, G. A., Mathers, C. D., Bonita, R., Rehm, J., Kruk, M. E., et al. (2018). NCD Countdown 2030: Worldwide trends in non-communicable disease mortality and progress towards Sustainable Development Goal target 3.4. *The Lancet, 392*(10152), 1072–1088.

Bull, F. C., Al-Ansari, S. S., Biddle, S., Borodulin, K., Buman, M. P., Cardon, G., et al. (2020). World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine, 54*(24), 1451–1462.

Ding, D., Lawson, K. D., Kolbe-Alexander, T. L., Finkelstein, E. A., Katzmarzyk, P. T., van Mechelen, W., & Pratt, M. (2016). The economic burden of physical inactivity: A global analysis of major non-communicable diseases. *The Lancet, 388*(10051), 1311–1324.

GBD 2019 Diseases and Injuries Collaborators. (2020). Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019. *The Lancet, 396*(10258), 1204–1222.

Guthold, R., Stevens, G. A., Riley, L. M., & Bull, F. C. (2018). Worldwide trends in insufficient physical activity from 2001 to 2016. *The Lancet Global Health, 6*(10), e1077–e1086.

Hallal, P. C., Andersen, L. B., Bull, F. C., Guthold, R., Haskell, W., & Ekelund, U. (2012). Global physical activity levels: Surveillance progress, pitfalls, and prospects. *The Lancet, 380*(9838), 247–257.

Hruby, A., & Hu, F. B. (2015). The epidemiology of obesity: A big picture. *Pharmacoeconomics, 33*, 673–689.

Kivimäki, M., Kuosma, E., Ferrie, J. E., Luukkonen, R., Nyberg, S. T., Alfredsson, L., et al. (2017). Overweight, obesity, and risk of cardiometabolic multimorbidity. *The Lancet Diabetes & Endocrinology, 5*(4), 253–263.

CC BY

ISSN 2674-8584

Received: 17/07/2026 - **Accepted:** 21/08/2026
Vol: 08.08 - August
DOI: 10.66104/zxqd3c29
Pages: 1-13

Lavie, C. J., Ozemek, C., Carbone, S., Katzmarzyk, P. T., & Blair, S. N. (2019). Sedentary behavior, exercise, and cardiovascular health. *Circulation Research, 124*(5), 799–815.

NCD Risk Factor Collaboration. (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016. *The Lancet, 390*(10113), 2627–2642.

Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., et al. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013. *The Lancet, 384*(9945), 766–781.

Pan American Health Organization. (2023). *Plan of action for the prevention and control of noncommunicable diseases*. Pan American Health Organization.

Popkin, B. M., Adair, L. S., & Ng, S. W. (2012). Global nutrition transition and the pandemic of obesity in developing countries. *Nutrition Reviews, 70*(1), 3–21.

Swinburn, B. A., Kraak, V. I., Allender, S., Atkins, V. J., Baker, P. I., Bogard, J. R., et al. (2019). The global syndemic of obesity, undernutrition, and climate change. *The Lancet, 393*(10173), 791–846.

World Health Organization. (2022). *Global status report on physical activity 2022*. World Health Organization.

CC BY